

**BULLETIN
OF
ENTOMOLOGICAL
RESEARCH**

VOL. 79

**ISSUED BY THE C·A·B INTERNATIONAL
INSTITUTE OF ENTOMOLOGY**

EDITOR: J. M. B. HARLEY

LONDON

**C·A·B INTERNATIONAL INSTITUTE OF ENTOMOLOGY
56, QUEEN'S GATE, SW7 5JR**

1989

C·A·B INTERNATIONAL INSTITUTE OF ENTOMOLOGY

Director

K. M. HARRIS, BSc, DSc, DipAgrSc, DTA, FIBiol

Editor

J. M. B. HARLEY, BSc

Headquarters and Publication Office and Library—56 Queen's Gate, London SW7 5JR, UK.

Telephone: 01-584-0067. Telegrams: ENTINFO LONDON SW7. BT GOLD/DIALCOM: 84: CAU006. Telex: 265871 MONREF G (Box CAU006). Fax: 01-581-1676.

Identification Service—c/o British Museum (Natural History), Cromwell Road, London SW7 5BD, UK.

The C·A·B International Institute of Entomology is one of 4 Institutes and 11 Bureaux that together form C·A·B International (CABI) a not-for-profit, inter-governmental organisation which exists to provide information, scientific and development services for agriculture and allied disciplines throughout the world.

The *Bulletin of Entomological Research* is a quarterly journal prepared by the Institute and published by C·A·B International.

Annual Subscription Rate 1989

| | |
|--|---------|
| To subscribers in countries that are members of CABI | £77·00 |
| To subscribers in countries that are not members of CABI | £103·00 |

Back Numbers

Back volumes of the *Bulletin of Entomological Research* are available at a discount.

Orders and enquiries concerning subscriptions and back numbers should be sent to

Marketing and Sales Division,
C·A·B INTERNATIONAL,
WALLINGFORD, OXON, OX10 8DE, UK.
TELEPHONE: WALLINGFORD (0491) 32111
TELEX: 847964 (COMAGG G)
FAX: (0491) 33508
TELECOM GOLD/INTERNATIONAL DIALCOM: 84: CAU001

[Note: Prices are subject to change without notice.]

© C·A·B International, 1989. All rights reserved. No part of this publication may be reproduced in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, without the prior permission of the copyright owner.

Disclaimer

The information contained herein, including any expression of opinion and any projection or forecast, has been obtained from or is based upon sources believed by us to be reliable but is not guaranteed as to accuracy or completeness. The information is supplied without obligation and on the understanding that any person who acts upon it or otherwise changes his/her position in reliance thereon does so entirely at his/her own risk.

C·A·B International does not accept responsibility for any trade advertisement included in this publication.

CONTENTS

PART 1 MARCH 1989

Original articles

| | PAGE |
|---|------|
| QUICKE, D. L. J. & WHARTON, R. A. <i>Myosoma nyanzaensis</i> sp. n. (Hymenoptera: Braconidae) parasitic on the stem-borer pest <i>Chilo partellus</i> (Swinhoe) (Lepidoptera: Pyralidae) in Kenya | 1 |
| ROBERTSON, H. G. & HOFFMANN, J. H. Mortality and life-tables of <i>Cactoblastis cactorum</i> (Berg) (Lepidoptera: Pyralidae) compared on two host-plant species | 7 |
| TYNDALE-BISCOE, M. The influence of adult size and protein diet on the human-oriented behaviour of the bush fly, <i>Musca vetustissima</i> Walker (Diptera: Muscidae) | 19 |
| BARNES, B. N. & PRINGLE, K. L. Oviposition by the banded fruit weevil, <i>Phylloctenus callosus</i> (Schoenherr) (Coleoptera: Curculionidae), in deciduous fruit orchards in South Africa | 31 |
| KAPATSA, G., SPATES, G. E., SHEFFIELD, C. L. & DELOACH, J. R. Comparison of lipid-free haemoglobin and stroma-contaminated haemoglobin diets for adults of <i>Stomoxys calcitrans</i> (L.) (Diptera: Muscidae) | 41 |
| LOPEZ, E. R. & VAN DRIESCHE, R. G. Direct measurement of host and parasitoid recruitment for assessment of total losses due to parasitism in a continuously breeding species, the cabbage aphid, <i>Brevicoryne brassicae</i> (L.) (Hemiptera: Aphididae) | 47 |
| ATKINSON, P. R. & CARNEGIE, A. J. M. Population dynamics of the sugarcane borer, <i>Eldana saccharina</i> Walker (Lepidoptera: Pyralidae), in Natal, South Africa | 61 |
| GREEN, C. H. The use of two-coloured screens for catching <i>Glossina palpalis palpalis</i> (Robineau-Desvoidy) (Diptera: Glossinidae) | 81 |
| KABISSA, J. C. B. Evaluation of damage thresholds for insecticidal control of <i>Helicoverpa armigera</i> (Hübner) (Lepidoptera: Noctuidae) on cotton in eastern Tanzania | 95 |
| TORR, S. J., PARKER, A. G. & LEIGH-BROWNE, G. The responses of <i>Glossina pallidipes</i> Austen (Diptera: Glossinidae) to odour-baited traps and targets in Somalia | 99 |
| PÄTS, P. & WIKTELius, S. Tethered flight of <i>Chilo partellus</i> (Swinhoe) (Lepidoptera: Pyralidae) | 109 |
| GOLD, C. S., ALTIERI, M. A. & BELLOTTI, A. C. The effects of intercropping and mixed varieties of predators and parasitoids of cassava whiteflies (Hemiptera: Aleyrodidae) in Colombia | 115 |
| HEMINGWAY, J., BODDINGTON, R. G., HARRIS, J. & DUNBAR, S. J. Mechanisms of insecticide resistance in <i>Aedes aegypti</i> (L.) (Diptera: Culicidae) from Puerto Rico | 123 |
| AUSTIN, A. D. & DANGERFIELD, P. C. The taxonomy of New World microgastrine braconids (Hymenoptera) parasitic on <i>Diatraea</i> spp. (Lepidoptera: Pyralidae) | 131 |
| FITT, G. P., ZALUCKI, M. P. & TWINE, P. Temporal and spatial patterns in pheromone-trap catches of <i>Helicoverpa</i> spp. (Lepidoptera: Noctuidae) in cotton-growing areas of Australia | 145 |
| LIEBREGTS, W. J. M. M., SANDS, D. P. A. & BOURNE, A. S. Population studies and biological control of <i>Pseudaulacaspis pentagona</i> (Targioni-Tozzetti) (Hemiptera: Diaspididae) on passion fruit in Western Samoa | 163 |

Review article

| | PAGE |
|--|------|
| COWIE, R. H., LOGAN, J. W. M. & WOOD, T. G. Termite (Isoptera) damage and control in tropical forestry with special reference to Africa and Indo-Malaysia: a review | 173 |
| Original articles | |
| MCGEACHIE, W. J. The effects of moonlight illuminance, temperature and wind speed on light-trap catches of moths | 185 |
| MALCOLM, C. A. & BODDINGTON, R. G. Malathion resistance conferred by a carboxylesterase in <i>Anopheles culicifacies</i> Giles (species B) (Diptera: Culicidae) | 193 |
| TAHHAN, O. & VAN EMDEN, H. F. Biology of <i>Bruchus dentipes</i> Baudi (Coleoptera: Bruchidae) on <i>Vicia faba</i> and a method to obtain gravid females during the imaginal quiescence period | 201 |
| TAHHAN, O. & VAN EMDEN, H. F. Resistance of faba bean, <i>Vicia faba</i> , to <i>Bruchus dentipes</i> Baudi (Coleoptera: Bruchidae) | 211 |
| WAY, M. J., CAMELL, M. E., BOLTON, B. & KANAGARATNAM, P. Ants (Hymenoptera: Formicidae) as egg predators of coconut pests, especially in relation to biological control of the coconut caterpillar, <i>Opisina arenosella</i> Walker (Lepidoptera: Xyloryctidae), in Sri Lanka | 219 |
| BRAUN, A. R., GUERRERO, J. M., BELLOTTI, A. C. & WILSON, L. T. Within-plant distribution of <i>Mononychellus tanajoa</i> (Bondar) (Acari: Tetranychidae) on cassava: effect of clone and predation on aggregation | 235 |
| ALLSOPP, P. G. & BULL, R. M. Spatial patterns and sequential sampling plans for melolonthine larvae (Coleoptera: Scarabaeidae) in southern Queensland sugarcane | 251 |
| PEGRAM, R. G., ZIVKOVIC, D. & JONGEJAN, F. Ticks (Acari: Ixodoidea) of the Yemen Arab Republic. II. The <i>Rhipicephalus sanguineus</i> (Latreille) group | 259 |
| WILSON, A. G. L. & MORTON, R. Some factors affecting the reliability of pheromone traps for measurement of the relative abundance of <i>Helicoverpa punctigera</i> (Wallengren) and <i>H. armigera</i> (Hübner) (Lepidoptera: Noctuidae) | 265 |
| JACKAI, L. E. N. A laboratory procedure for rearing the cowpea coreid, <i>Clavigralla tomentosicollis</i> Stål (Hemiptera), using dry cowpea seeds | 275 |
| WALKER, A. K. Synonymy of two <i>Apanteles</i> species (Hymenoptera: Braconidae) parasitizing the coconut flat moth, <i>Agonoxena argaula</i> Meyrick (Lepidoptera: Agonoxenidae) | 283 |
| ASCHE, M. & WILSON, M. R. The three taro planthoppers: species recognition in <i>Tarophagus</i> (Hemiptera: Delphacidae) | 285 |
| HOSSAIN, M. I., CURTIS, C. F. & HEEKIN, J. P. Assays of permethrin-impregnated fabrics and bioassays with mosquitoes (Diptera: Culicidae) | 299 |
| PADGHAM, D. E., WOODHEAD, S. & RAPUSAS, H. R. Feeding responses of the brown planthopper, <i>Nilaparvata lugens</i> (Stål) (Hemiptera: Delphacidae), to resistant and susceptible host-plants | 309 |
| HALL, M. J. R. & LANGLEY, P. A. The responses of individual males in an isolated population of <i>Glossina morsitans morsitans</i> Westwood (Diptera: Glossinidae) to pheromone-baited decoy 'females' | 319 |
| WALL, R. Sexual responses of males of <i>Glossina morsitans morsitans</i> Westwood and <i>G. pallidipes</i> Austin (Diptera: Glossinidae) to traps and targets in the field | 335 |

Review article

LEATHER, S. R., WALTERS, K. F. A. & DIXON, A. F. G. Factors determining the pest status of the bird cherry-oat aphid, *Rhopalosiphum padi* (L.) (Hemiptera: Aphididae), in Europe: a study and review 345

Original articles

AMIN, A. M. & HEMINGWAY, J. Preliminary investigation of the mechanisms of DDT and pyrethroid resistance in *Culex quinquefasciatus* Say (Diptera: Culicidae) from Saudi Arabia 361

NOYES, J. S. *Copidosoma graminis* sp.n. (Hymenoptera: Encyrtidae), a polyembryonic parasitoid of armyworms (Lepidoptera: Noctuidae) in Australia 367

ZHOU, X., CARTER, N. & MUMFORD, J. A simulation model describing the population dynamics and damage potential of the rose grain aphid, *Metopolophium dirhodum* (Walker) (Hemiptera: Aphididae), in the UK 373

BAILEY, P. T. The millipede parasitoid *Pelidnoptera nigripennis* (F.) (Diptera: Sciomyzidae) for the biological control of the millipede *Ommatoiulus moreleti* (Lucas) (Diplopoda: Julidae) in Australia 381

ANSART, A. K., VAN EMDEN, H. F. & SINGH, S. R. Graft-transmissibility of resistance to cowpea aphid, *Aphis craccivora* Koch (Hemiptera: Aphididae), in six highly antibiotic cowpea varieties 393

WHARTON, R. A., SMITH, J. W., JR, QUICKE, D. L. J. & BROWNING, H. W. Two new species of *Digonogastra* Viereck (Hymenoptera: Braconidae) parasitic on Neotropical pyralid borers (Lepidoptera) in maize, sorghum and sugarcane. 401

SATTLER, K. & STRIDE, A. B. A new species of *Hypatima* Hübner (Lepidoptera: Gelechiidae) injurious to mango trees in East Africa 411

BALLARD, J. W. O. Factors influencing silhouette-trap captures of the blackfly *Austrosimulium bancrofti* (Taylor) (Diptera: Simuliidae) in the Australian Capital Territory 421

LANGLEY, P. A. Laboratory colonization of the tsetse fly *Glossina pallidipes* Austen (Diptera: Glossinidae) using an *in vitro* feeding method 429

COOK, A. G. & PERFECT, T. J. Population dynamics of three leafhopper vectors of rice tungro viruses, *Nephrotettix virescens* (Distant), *N. nigropictus* (Stål) and *Recilia dorsalis* (Motschulsky) (Hemiptera: Cicadellidae), in farmers' fields in the Philippines 437

WHITE, I. M. & MARQUARDT, K. A revision of the genus *Chaetorellia* Hendel (Diptera: Tephritidae) including a new species associated with spotted knapweed, *Centaurea maculosa* Lam. (Asteraceae) 453

ATKINSON, P. R. & NUSS, K. L. Associations between host-plant nitrogen and infestations of the sugarcane borer, *Eldana saccharina* Walker (Lepidoptera: Pyralidae) 489

NEWTON, P. J. Combinations of applications of a chitin synthesis inhibitor and inundative releases of egg parasitoids against the false codling moth, *Cryptophlebia leucotreta* (Meyrick) (Lepidoptera: Tortricidae), on citrus 507

DOKU, C. & BRADY, J. Landing site preferences of *Glossina morsitans morsitans* Westwood (Diptera: Glossinidae) in the laboratory: avoidance of horizontal features? 521

Original articles

| | PAGE |
|--|------|
| WRIGHT, A. D., BOLAND, N. P. & KERR, J. D. Changes in leaf hardness and moisture content of water hyacinth, <i>Eichhornia crassipes</i> , following feeding by <i>Neochetina eichhorniae</i> Warner (Coleoptera: Curculionidae) | 529 |
| VAN DER LINDE, T. C. DE K., HEWITT, P. H., VAN DER WESTHUIZEN, M. C. & MITCHELL, J. The use of ^{131}I , ^{125}I and aggressive behaviour to determine the foraging area of <i>Hodotermes mossambicus</i> (Hagen) (Isoptera: Hodotermitidae) | 537 |
| DIKE, M. C. A key for the identification of Afrotropical species of the shoot-fly subgenus <i>Atherigona</i> of <i>Atherigona</i> (Diptera: Muscidae), with a description of some new species from Africa | 545 |
| WHEATLEY, A. R. D., WIGHTMAN, J. A., WILLIAMS, J. H. & WHEATLEY, S. J. The influence of drought stress on the distribution of insects on four groundnut genotypes grown near Hyderabad, India | 567 |
| NEUENSCHWANDER, P., HAMMOND, W. N. O., GUTIERREZ, A. P., CUDJOE, A. R., ADJAKLOE, R., BAUMGÄRTNER, J. U. & REGEV, U. Impact assessment of the biological control of the cassava mealybug, <i>Phenacoccus manihoti</i> Matile-Ferrero (Hemiptera: Pseudococcidae), by the introduced parasitoid <i>Epidinocarsis lopezi</i> (De Santis) (Hymenoptera: Encyrtidae) | 579 |
| JACKAI, L. E. N. & OGHIAKHE, S. Pod wall trichomes and resistance of two wild cowpea, <i>Vigna vexillata</i> , accessions to <i>Maruca testulalis</i> (Geyer) (Lepidoptera: Pyralidae) and <i>Clavigralla tomentosicollis</i> Stål (Hemiptera: Coreidae) | 595 |
| MARTIN, J. H. Identification, occurrence and pest status of <i>Toxoptera odinae</i> (van der Goot) (Hemiptera: Aphididae) in Africa | 607 |
| JOHNSTONE, D. R., COOPER, J. F., DOBSON, H. M. & TURNER, C. R. The collection of aerosol droplets by resting tsetse flies, <i>Glossina morsitans</i> Westwood (Diptera: Glossinidae) | 613 |
| SULAIMAN, S., SOHADI, A. R. & JEFFERY, J. Human helminth parasite burdens on cyclorrhaphan flies (Diptera) trapped at an aboriginal settlement in Malaysia | 625 |
| DUNCAN, F. D. & HEWITT, P. H. Observations on the foraging behaviour of the harvester termite, <i>Hodotermes mossambicus</i> (Hagen) (Isoptera: Hodotermitidae) | 631 |
| WILLIAMS, D. J. A new genus and two new species of mealybugs (Hemiptera: Pseudococcidae) from South Africa on bulbs and corms imported into Great Britain | 643 |
| ADAIR, R. F. & SCOTT, J. K. The life-history and host specificity of <i>Comostolopsis germana</i> Prout (Lepidoptera: Geometridae), a biological control agent of <i>Chrysanthemoides monilifera</i> (Compositae) | 649 |
| HEMINGWAY, J., CALLAGHAN, A. & KURTAK, D. C. Temephos resistance in <i>Simulium damnosum</i> Theobald (Diptera: Simuliidae): a comparative study between larvae and adults of the forest and savanna strains of this species complex .. | 659 |
| AGRICOLA, U., AGOUNKÉ, D., FISCHER, H. U. & MOORE, D. The control of <i>Rastrococcus invadens</i> Williams (Hemiptera: Pseudococcidae) in Togo by the introduction of <i>Gyranusoidea tebygi</i> Noyes (Hymenoptera: Encyrtidae) .. | 671 |
| EL-SAYED, S., HEMINGWAY, J. & LANE, R. P. Susceptibility baselines for DDT metabolism and related enzyme systems in the sandfly <i>Phlebotomus papatasii</i> (Scopoli) (Diptera: Psychodidae) | 679 |
| THOMSON, M. C., DAVIES, J. B. & WILSON, M. D. A portable allozyme electrophoresis kit used to identify members of the <i>Simulium damnosum</i> Theobald complex (Diptera: Simuliidae) in the field | 685 |
| HOBBS, S. E. & WOLF, W. W. An airborne radar technique for studying insect migration | 693 |

DATES OF PUBLICATION IN PARTS

| | |
|---------------------------|-------------------|
| PART 1, pp. 1-171 | 16 March 1989 |
| PART 2, pp. 173-343 | 23 May 1989 |
| PART 3, pp. 345-528 | 13 September 1989 |
| PART 4, pp. 529-720 | 30 November 1989 |

ERRATA (VOL. 79)

p. 115, title line 1, for 'of predators' read 'on predators'.

p. 116, line 24 and p. 120, Table IV & line 22, for 'Amitus aleurodinus' read 'Amitus aleurodinis'.

p. 120, Table IV, title line 2, for 'aleurodiphaga' read 'aleyrodiphaga'.

pp. 235-249 except p. 235 line 20, for 'tanajoa' read 'progresivus'.

ADDENDA (VOL. 79)

pp. 285-298, article by M. Asche & M. R. Wilson.

By an oversight only one species of *Tarophagus*, *T. colocasiae* (Matsumara) was recorded from Hawaii. Further specimens have now been examined in the B.P. Bishop Museum, Honolulu and in the Dept. of Entomology, University of Hawaii at Manoa. We now find that *T. proserpina* (Kirkaldy) is also found in Hawaii and, indeed, is more common than *T. colocasiae*. Specimens are recorded from several localities on Oahu Island. The following additions/corrections should be made to the paper as follows:

p. 290, 'T. proserpina, Material examined' add 'Hawaii Islands (Oahu)'.

p. 295, Table I, add 'x' for *T. proserpina* at Hawaii.

p. 296, Fig. 35 is revised and an error in the legend corrected as below




Fig. 35.—Distribution map of *Tarophagus* species.

p. 296, Table II, add 'at Hawaii under probable species of *Tarophagus*' add 'or *proserpina*'.
p. 296, Table III, revised version is given below:

| To | <i>Tarophagus</i> sp. | From | <i>Tarophagus</i> sp. | Result |
|----------------------|--|-------------|--|--------|
| Tahiti | <i>proserpina</i> | Hawaii | <i>colocasiae</i> or <i>proserpina</i> | - |
| Guam | <i>colocasiae</i> | Hawaii | <i>colocasiae</i> or <i>proserpina</i> | + |
| Hawaii | <i>colocasiae</i> or <i>proserpina</i> | Philippines | <i>colocasiae</i> or <i>persephone</i> | + |
| Ponape (Caroline Is) | <i>colocasiae</i> | Guam | <i>colocasiae</i> | + |
| Western Samoa | <i>proserpina</i> | Hawaii | <i>colocasiae</i> or <i>proserpina</i> | + |
| Solomon Is | <i>persephone</i> or <i>colocasiae</i> | Sikaiana | <i>persephone</i> or <i>colocasiae</i> | + |
| | " | Samoa | <i>colocasiae</i> | - |
| Tuvalu | <i>proserpina</i> | Fiji | <i>proserpina</i> | - |